

Progress Toward a Large-Scale Synthesis of Molnupiravir (MK-4482, EIDD-2801) from Cytidine

Grace P. Ahlqvist, Catherine P. McGeough, Chris Senanayake, Joseph D. Armstrong, Ajay Yadaw, Sarabindu Roy, Saeed Ahmad, David R. Snead, and Timothy F. Jamison*



Cite This: *ACS Omega* 2021, 6, 10396–10402



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Molnupiravir (MK-4482, EIDD-2801) is a promising orally bioavailable drug candidate for the treatment of COVID-19. Herein, we describe a supply-centered and chromatography-free synthesis of molnupiravir from cytidine, consisting of two steps: a selective enzymatic acylation followed by transamination to yield the final drug product. Both steps have been successfully performed on a decagram scale: the first step at 200 g and the second step at 80 g. Overall, molnupiravir has been obtained in a 41% overall isolated yield compared to a maximum 17% isolated yield in the patented route. This route provides many advantages to the initial route described in the patent literature and would decrease the cost of this pharmaceutical should it prove safe and efficacious in ongoing clinical trials.



- Two-step, lower-cost synthesis of potential COVID-19 therapeutic
- Selective acylation using commercial solid-supported enzyme
- Chromatography-free purification by recrystallization

INTRODUCTION

Molnupiravir (MK-4482, EIDD-2801) is in development by Merck after licensing from Ridgeback Biopharmaceuticals as an orally dosed antiviral for the treatment of COVID-19.¹ Animal studies have shown successful inhibition of SARS-CoV-2² as well as prevention of viral transmission.³ If shown to be safe and effective in ongoing clinical trials, this compound would be an important tool in the toolbox of physicians working to counter the effects of the SARS-CoV-2 virus pandemic. Drug availability, however, depends on the efficient and cost-effective synthesis of the drug molecule to ensure broad global access to this potentially valuable medication. The original synthetic route was disclosed by Emory University in 2019; this route made molnupiravir in five steps with uridine (**1**) as the starting material.⁴ Since the yield for the final two steps is not reported, this route has a 17% maximum overall yield (Scheme 1).

We previously demonstrated the potential of a two-step route from cytidine (**8**) to molnupiravir (**7**), which has many advantages over the previously patented route including cost and the overall yield (Figure 1).⁵

However, several challenges prevent the previously reported route from being implemented at the manufacturing scale. Most notably, both the intermediate and the final active pharmaceutical ingredient (API) were purified by column chromatography, so we sought to develop alternate workup and crystallization procedures to provide a pure material without chromatographic purification. Other opportunities for improvement included examining the surprisingly high cost of acetone oxime and refining process parameters such as reaction concentration, catalyst loading, and stoichiometry. Thus, we

set out to re-examine this route to enable this process to be run at an increased scale.

RESULTS AND DISCUSSION

The oxime ester acylating agent (**9**) proved uniquely suited to selective enzyme acylation (Supporting Information (SI) Section 2.1). Unfortunately, the acetone oxime used in the synthesis of the acylating agent proved to be a major cost driver of our initial route investigation (SI, Section 1).

Surprised by the high cost of such a simple material, we sought to determine the cost of synthesizing this material ourselves. Gratifyingly, we found that a reported procedure⁶ for acetone oxime synthesis provided 64% yield on our first attempt (Scheme 2a). The use of methyl *tert*-butyl ether (MTBE) to replace diethyl ether in the workup produced a yield of 71%. This method was repeated on a 500 g scale with NaOH as the base, and 73% yield was observed. We anticipate a significant reduction in the cost of acetone oxime prepared by this procedure compared to current commercial sources.

With a cheaper preparation of acetone oxime in hand, we also sought to improve the synthesis of oxime ester **9**. Our previous synthesis (Scheme 2b) used a slight excess of isobutyric acid chloride and triethylamine and 50 V of dichloromethane (DCM) as the solvent. We found that we



Received: February 10, 2021

Accepted: March 30, 2021

Published: April 8, 2021




Scheme 1. Synthetic Route of Molnupiravir Disclosed by Emory University in 2019



This work	Previous work
No column chromatography purifications	All column chromatography purifications
Step 1: 200 g scale	Step 1: 5 g scale
Step 2: 80 g scale	Step 2: 1 g scale

Figure 1. This work (green) compared to the previous work (blue).

Scheme 2. (a) Synthesis of Acetone Oxime (11); (b) Synthesis of Acylating Agent 9 Using Acetone Oxime (11) and Isobutyryl Chloride (13)



could increase the throughput of the reaction by decreasing the solvent to 30 V, as well as increasing triethylamine addition from 1.1 to 1.2 equiv. These small changes afforded a 99% corrected yield on a 35 g scale. This reaction was also scaled up significantly, to 200 g, with a 90% corrected yield. No purification was necessary as the crude material was sufficient for use in the next step (*vide infra*). We hypothesize that this reaction could be further improved by replacing dichloromethane with a less hazardous solvent and that the reaction scale could be increased even further.

Streamlined preparation of acylating agent 9 facilitated investigation of the enzymatic acylation of cytidine (Figure 1, step 1). We first sought to investigate environmentally

preferable solvents to substitute for 1,4-dioxane (dioxane).⁷ Unfortunately, extensive solvent screening (Figure 2, additional data in SI, Section 2.2) did not reveal any other solvent that worked as well as dioxane for the reaction.

Cyclopentanone appeared promising based on our initial screening, and its high boiling point (131 °C) opened the possibility for an increased reaction temperature that we hoped would accelerate conversion. Unfortunately, higher temperatures led to lower conversion (SI, Section 2.3), possibly due to the release of the enzymes from the polymer beads⁸ or degradation of the product. The lower conversion with cyclopentanone led us to return to dioxane as the optimal solvent. Finally, we did a cursory investigation to ascertain




Figure 2. Various solvents screened for the enzymatic acylation of cytidine (MeCN = acetonitrile, CPME = cyclopentyl methyl ether, 2-MeTHF = 2-methyl tetrahydrofuran).




Figure 3. Assessment of solvent volume versus enzyme loading. See SI, Section 2.7, for additional details.

whether a lower temperature or concentration could benefit the reaction; we found that lowering the temperature to 40 °C lowered conversion (SI, Section 2.4). We then screened other enzymes in dioxane and additional solvents, but the originally identified Novozym-435 (N435) was the most efficient and selective enzyme for the desired 5'-O-acylation (SI, Section 2.5). It was determined that a lower stoichiometry of the acylating agent (4.0 vs 5.0 equiv) could be used (SI, Section 2.6); while the yield decreased slightly, we believe the savings in cost and purity offset the decrease in the yield. Thus, 4.0 equiv of oxime ester and 60 °C were selected as the optimal conditions.

Hoping to decrease further the amount of the solvent needed for the reaction, a two-factor, four-level full factorial screening of 20–100 solvent volumes and 50–400 weight percent enzyme loading were performed (Figure 3).

We hoped that increased enzyme loading would allow us to decrease solvent usage, which would in turn improve the reaction throughput and process mass intensity (PMI). Unfortunately, our screening revealed that increasing concentration significantly hindered conversion; we hypothesize that this effect is due to the limited solubility of cytidine in dioxane. Ultimately, we chose conditions using 4.0 equiv of oxime ester, 150 wt % N435, and 50 V of dioxane as the solvent, due to the

highest distribution of product 10 relative to remaining starting material 8 and diacylated product 14 (SI, Section 3.3).

Additionally, the breadth of data on regenerating and recycling N435, including repeated uses in organic solvents,^{8,9a-d} suggests that N435 could be recycled for this reaction as well. Reuse of the enzyme would further decrease the raw material costs associated with this route.

Upon scaling up the reaction, we noticed a slowed reaction rate, from 24 h to completion at a small scale to approximately 40 h at a 19 g scale (SI, Section 2.8). We believe that this change is related to differences in the stirring method used at different scales: we stirred using magnetic stir bars at 750 rpm for scales up to 1 g, while larger-scale reactions used an overhead stirrer set to 260 rpm. Using this same setup, the reaction was scaled 10-fold to 200 g with no change in performance.

Throughout the reaction optimization, the major impurity observed was the diacylated product (14), the structure of which was confirmed as the N-acylated product via ¹⁵N-¹H 2-D NMR spectroscopy (SI, Section 2.9). However, we also saw a consistent unknown impurity in the high-performance liquid chromatography (HPLC), constituting 5–10 LC area percent (SI, Section 2.10). We noticed that this impurity increased in proportion with enzyme loading, leading us to believe that the


Table 1. Cytidine Acylation Summary

8 Scale (g)	Solvent vol.	enzyme wt. %	Oxime ester 9 equiv	Reaction Time (hours)	Product 10 IY	Purity ^a
20	40 V	150	4.0	38	60%	>99%
30	50 V	150	4.0	40	65%	>99%
100	52.5 V	150	3.2	43	68%	98%
100	60 V	150	3.2	43	70%	>99% ^c
19	50 V	150	4.0	42	66%	>99%
200	52.5 V	150	4.0	43	70%	96%

^aPurity determined by qNMR unless noted otherwise; ^bAdditional details for purification in experimental section; ^cPurity determined by HPLC (at 280 nm)

Workup A	Add acetone (10 V), reflux (30 min), cool then filter, rinse with MTBE (2V)
Workup B	Triturate with MTBE (10 V), add acetone (13 V), reflux, cool, then filter
Workup C	Add water (20 V): MTBE (20 V) to solid, extract MTBE layer with water (2 x), remove water from combined water extracts, add acetone (13 V), reflux, cool, then filter

Table 2. Development of Conditions for the Transamination of Compound 10 to Molnupiravir (7)



entry	scale of 10 (g)	(NH ₂ OH) ₂ H ₂ SO ₄ equiv	time (h)	7 crude purity (%) ^a	10 LC area % ^b	7 LC area % ^b	15 LC area % ^b
1	1	4.5	22	73	9	89	2
2	1	3	22	85	8	90	2
3	1	2	22	84	9	89	2
4	5	3	22	97	6	92	2

^aPurity determined by qNMR. ^bDetermined by LC-MS at 280 nm.

impurity was leached from the enzyme beads by dioxane. Indeed, a control reaction confirmed that the impurity formed simply from stirring the enzyme beads in dioxane at 60 °C overnight and was not related to either the cytidine or the acylating agent (SI, Section 2.9). Isolation of this impurity yielded a yellow gel that did not ionize on liquid chromatography-mass spectrometry (LC-MS). We hypothesize that this material is a macromolecular compound leached from the Lewatit VP OC 1600 solid support; solubilization of the support in studies of N435 catalysis has been reported for some solvents.⁸ Fortunately, simply rinsing the enzymes with dioxane before use eliminated this impurity (SI, Section 2.9). Thus, the main impurities remaining in the reaction were the diacylated product 14, unreacted cytidine 8, and excess acylating agent 9. Allowing this reaction to cool to room temperature (RT, 20 °C) and then filtering enabled the removal of the enzyme beads as well as the unreacted starting material (SI, Section 3.2); this was due to cytidine's insolubility in room-temperature dioxane.

The main challenge for purification was therefore removing the excess acylating agent and the diacylated product from the desired product 10. Upon completion of the reaction, the

remaining cytidine starting material as well as the enzyme beads were filtered out and then dioxane was removed via rotary evaporation. We then determined three methods (A, B, and C) for the purification of 10 from compounds 14 and 9 (Table 1). Methods A and B were successful for the purification of desired product 3 up to a purity of >99% on smaller scales (10–100 g), while purification C was successful on up to 200 g scale.

The previous report of the transamination of intermediate 10, although high yielding, was obtained using column chromatography, which would not be viable on the scale.⁵ Previously, 70% aq isopropanol was used as the solvent for this reaction; however, we chose to replace this with 70% aq 1-butanol solution. We hypothesized that the decreased solubility of butanol in water (as compared to the solubility of isopropanol in water) would decrease the amount of hydroxylamine salts present in the organic layer after reaction, thus leading to a more facile purification. The optimization of this transamination reaction began by using 4.5 equiv of hydroxylamine sulfate relative to the starting material 10,⁵ which displayed a good conversion of 10 to the desired product 7 after 22 h (SI, Section 4.1). In addition to the

Table 3. Scale-Up of the Transamination of 10 to 7 as well as the Isolated Yields and Purity of the Isolated Material

entry	scale of 10 (g)	(NH ₂ OH) ₂ H ₂ SO ₄ equiv	time (h)	7 crude purity (%) ^a	10 LC area % ^a	7 LC area % ^a	15 LC area % ^a	7 isolated yield (%)	purity of isolated 7 (%) ^b
1	10	3	24	92	4	94	2	48	>99
2	20	3	28	93	2	96	2	49	>99
3	80	3.2	40	80	4	91	2	58	97

^aBased on LC-MS area percent at 280 nm. ^bPurity determined by qNMR.

desired product formation, the side product *N*-hydroxycytidine (NHC, 15) was formed in this reaction in small quantities. We found that decreasing the stoichiometry of hydroxylamine sulfate from 4.5 equiv to 2 or 3 equiv led to very similar reaction conversion (Table 2 and SI, Section 4.1) and increased the purity of the crude reaction mixture after workup, which simply involved separating the organic and aqueous layers and then removing the 1-butanol from the organic layer (crude purity, Table 2).

After the development of successful small-scale conditions (5 g), the next goals were to scale up this reaction to 80 g and to develop recrystallization conditions for the crude reaction mixture. We found that increasing the scale from 5 to 10 to 20 g did not significantly affect the reaction time. However, increasing the scale to 80 g required an increased reaction time to 40 h for a similar conversion (Table 3).

We found that the recrystallization purification of this reaction on a 80 g scale worked the same as on a 1 g scale, by simply separating the aqueous and organic layers. To increase the purity of the final compound from 80 to 90% after this initial purification, recrystallization conditions were developed. Due to the difference in solubility of compound 7 in cold versus hot water (Table 4), water was chosen as the

Table 4. Qualitative Solubility Studies of Molnupiravir in Water at Various Solvent Volumes and Temperatures

		Solution Temperature					
		5 °C	25 °C	40 °C	65 °C	80 °C	95 °C
Water Volume	20						
	15						
	10						
	5						
	2						
	1						
	0.5						

	Soluble
	Partially soluble
	Not soluble

recrystallization solvent. Thus, it was determined that after the solvent was removed from the organic layer, 7 could be recrystallized in greater than 99% purity from heating to 65 °C and slowly cooling in water (2 V). After the initial recrystallization, the remainder of 7 remained in the aqueous filtrate. We anticipate that further recrystallizations of the filtrate would produce additional molnupiravir.

CONCLUSIONS

In conclusion, we have developed an increased scale, chromatography-free, two-step route from cytidine to molnupiravir with an overall isolated yield of 41% and lower cost as compared to the originally patented route³ (SI, Section 1), with further cost decreases possible if solvent and enzyme recycling is employed. We hope that further scale-up of this route will enable affordable, efficient preparation of this potentially crucial pharmaceutical for the global fight against COVID-19.

METHODS

General Procedures. For all compounds, ¹H and ¹³C NMR spectra were recorded on Bruker Avance III spectrometers (400, 500, or 600 MHz). Chemical shifts were measured relative to the residual solvent resonance for ¹H and ¹³C NMR (CDCl₃ = 7.26 ppm for ¹H and 77.2 ppm for ¹³C, DMSO-*d*₆ = 2.50 ppm for ¹H, and 39.2 ppm for ¹³C). Coupling constants *J* are reported in hertz (Hz). The following abbreviations were used to designate signal multiplicity: s, singlet; d, doublet; t, triplet; hept, heptet; dd, doublet of doublet; and m, multiplet. Reactions were monitored by HPLC (Agilent 1260 Infinity II LC) or LC-MS (Agilent Technologies InfinityLab LC/MSD). Unless noted otherwise, reactions involving air-sensitive reagents and/or requiring anhydrous conditions were performed under a nitrogen or argon atmosphere with glassware oven-dried at 140 °C. Reactions requiring mechanical stirring were stirred using a Heidolph RZR 2020 overhead stirring apparatus. Purity was assessed by quantitative NMR (qNMR) spectroscopy with benzyl benzoate or dimethyl sulfone (Sigma-Aldrich Trace-CERT grade) as the reference standard. Reagents and solvents were purchased from Aldrich Chemical Company, Fisher Scientific, Alfa Aesar, Acros Organics, Oakwood, or TCI. Unless otherwise noted, solid reagents were used without further purification. Methylene chloride (DCM) and dioxane were taken from a solid-sorbant Solvent Dispensing System purchased from Pure Process Technologies. Other solvents were purchased in anhydrous grades and used as received.

Acetone Oxime (11). 11 g Scale. To a solution of acetone (11.0 mL, 150 mmol, 1.0 equiv) and hydroxylamine hydrochloride (15.6 g, 225 mmol, 1.5 equiv) in HPLC-grade H₂O (300 mL) at RT was added Na₂CO₃ (28.6 g, 270 mmol, 1.8 equiv) in portions. The reaction mass was stirred for 19 h at RT, after which it was extracted with Et₂O (5 × 80 mL). The combined organic extracts were washed with saturated brine (40 mL) and dried over MgSO₄. After filtration and solvent removal, the product was obtained as a white solid in 64% yield (7.051 g) and 98% purity (qNMR). ¹H NMR (400 MHz, CDCl₃) δ 8.33 (s, 1H), 1.90 (s, 3H), and 1.89 (s, 3H). ¹H and ¹³C NMR matched reported literature values.⁶ Repetition of the reaction with MTBE instead of Et₂O in the workup improved the yield to 71% (7.735 g).

500 g Scale. To a solution of hydroxylamine hydrochloride (500 g, 7.20 mol, 1.1 equiv) in H₂O (670 mL) at 0–5 °C was added a solution of NaOH (270 g, 6.75 mol, 1 equiv) in H₂O (670 mL) dropwise. The reaction mass was stirred for 30 min at 0–5 °C and acetone (500 mL, 6.75 mol, 1 equiv) was added. The reaction mass was stirred for a further 2 h at 0–5 °C, after which it was filtered and vacuum-dried for 45 min yielding 450 g of wet white solid. The solid was then dissolved in DCM (9.0 L, 20 V) and stirred for 15 min, after which the layers were separated. The DCM layer was dried with Na₂SO₄ and washed with DCM (2.25 L, 5 V), then dried, yielding a white solid in 73% yield (360 g). The material was carried forward without further purification.

Acetone Oxime O-Isobutyryl Ester (9). 35 g Scale. Acetone oxime (35.0 g, 478.9 mmol, 1.0 equiv) was dissolved in dichloromethane (1050 mL, 30 V) under an argon atmosphere with mechanical stirring and cooled to 0 °C using an ice/water bath. Isobutyryl chloride (55.2 mL, 526.7 mmol, 1.1 equiv) was slowly added, maintaining the solution temperature between 0 and 5 °C. Et₃N (31.46 mL, 574.7 mmol, 1.2 equiv) was added via a syringe pump at a rate of 2 mL/min, again maintaining the solution temperature below 5 °C. Et₃N addition caused the evolution of vapor and the formation of solid precipitates. The reaction mass was stirred for 20 h and allowed to warm to room temperature. The reaction mass was then washed with H₂O (2 × 350 mL), 5% w/w solution of NaHCO₃ (2 × 250 mL), H₂O (1 × 350 mL), 1 N aq HCl (2 × 250 mL), H₂O (1 × 350 mL), and saturated brine solution (1 × 100 mL), followed by drying over MgSO₄. After filtration and solvent removal, the product was obtained as a pale yellow liquid in 68.81 g yield (99% corrected for purity) and 99% purity (qNMR, benzyl benzoate). The material was carried forward without purification. ¹H NMR (400 MHz, CDCl₃) δ 2.66 (hept, J = 7.0 Hz, 1H), 2.05 (s, 3H), 1.99 (s, 3H), 1.24 (d, J = 7.0 Hz, 6H). ¹H and ¹³C NMR matched previously reported values.⁵

200 g Scale. Acetone oxime (200 g, 2.74 mol, 1.0 equiv) was dissolved in DCM (6 L) and cooled to 0 °C using an ice/water bath. Isobutyryl chloride (325 mL, 3.01 mol, 1.1 equiv) and Et₃N (241 mL, 3.28 mol, 1.2 equiv) were slowly added, maintaining the solution temperature between 0 and 5 °C. The reaction mass was stirred for 20 h and allowed to warm to room temperature. The reaction mass was then washed with H₂O (2 × 2 L), 5% w/w solution of NaHCO₃ (2 × 1.4 L), H₂O (1 × 1.4 L), 1 N aq HCl (2 × 1.4 L), H₂O (1 × 2 L), and saturated brine solution (1 × 600 mL), followed by drying over Na₂SO₄. After filtration and solvent removal, the product was obtained as a pale yellow liquid in 385 g yield (90% corrected for purity) and 92% purity (qNMR). The material was carried forward without purification.

5'-O-Isobutyryl Cytidine (10). 30 g Scale. Novozym-435 (45.0 g) was rinsed with 1,4-dioxane (2 × 100 mL) in a sintered glass funnel and dried under vacuum for 20 min. To an oven-dried 3 L three-neck RBF were added cytidine (30.0 g, 123.3 mmol), rinsed enzyme beads, 1,4-dioxane (1.5 L, 50 V), and crude acetone oxime O-isobutyryl ester (70.65 g, 493.4 mmol). The reaction mixture was heated to 64 °C using a heating mantle and stirred using a mechanical stirrer for 39 h, monitoring by HPLC (see SI for HPLC traces). The reaction mixture was then cooled to room temperature and filtered and washed with dioxane (3 V). The solvent was removed by rotary evaporation, yielding the crude product as an off-white foam (the crude material for this reaction was then purified via

purification method A, see SI, Section 3.1, for other purification procedures).

Purification A: Acetone (300 mL, 10 V) was added to the solid and then refluxed for 30 min. The suspension was allowed to cool to room temperature and then cooled to 5 °C (12 h). The white solid was filtered and rinsed with MTBE (2 V) and then allowed to dry at 40 °C (12 h) under vacuum to yield compound **10** (25.2 g, 65%, >99% purity by qNMR). ¹H NMR (500 MHz, DMSO) δ 7.58 (d, J = 7.4 Hz, 1H), 7.19 (d, J = 23.1 Hz, 2H), 5.76 (d, J = 3.7 Hz, 1H), 5.73 (d, J = 7.4 Hz, 1H), 5.40 (d, J = 5.2 Hz, 1H), 5.19 (d, J = 5.9 Hz, 1H), 4.28 (dd, J = 12.2, 3.2 Hz, 1H), 4.18 (dd, J = 12.1, 5.4 Hz, 1H), 3.99–3.94 (m, 2H), 3.93–3.87 (m, 1H), 2.58 (hept, J = 7.0 Hz, 1H), 1.11 (d, J = 7.0 Hz, 6H). ¹H and ¹³C NMR matched the previously reported values.⁵

200 g Scale. Novozym-435 (300 g, 150 wt %) was rinsed with 1,4-dioxane (4 L) and dried under vacuum for 30 min. Cytidine (200 g, 822 mmol, 1 equiv), rinsed enzyme beads, 1,4-dioxane (10.5 L, 52.5 V), and crude acetone oxime O-isobutyryl ester (471 g, 3.29 mol, 4 equiv). The reaction mixture was heated to 60 °C and stirred for 43 h. The reaction mixture was then cooled to room temperature, filtered, and washed with dioxane (3 V). The solvent was removed by distillation.

Purification C: MTBE (20 V) and H₂O (20 V) were then added. The layers were separated and the organic layer was washed twice with H₂O. The combined aqueous layers were distilled to yield a white solid. Acetone (13 V) was added and the slurry was heated to 60 °C for 1 h, cooled to room temperature, filtered, and dried, yielding 188 g (70% corrected) with a purity of 96.5% (qNMR).

Molnupiravir (7). 10 g Scale. 5'-O-Isobutyryl cytidine (10.0 g, 32.0 mmol, 1.0 equiv) was dissolved in 70% aq 1-butanol (112 mL of 1-butanol: 48 mL of water) and then hydroxylamine sulfate (15.8 g, 96.0 mmol, 3.0 equiv) was added. The mixture was stirred vigorously and heated to 78 °C for 24 h. The layers were then separated, and 1-butanol was removed from the organic layer via rotary evaporation to yield the solid white, crude material (5.2 g, 92% purity via qNMR). This crude material was then dissolved in water (2 V) and heated to 65 °C for 30 min (60 min for 80 g scale). After completely dissolved, the mixture was allowed to cool to room temperature and then to 5 °C without stirring. The solid was then filtered and washed with MTBE (2 V) to obtain molnupiravir (5.03 g, 48% yield, >99% purity by qNMR). ¹H NMR (500 MHz, DMSO) δ 10.00 (s, 1H), 9.53 (s, 1H), 6.83 (d, J = 8.2 Hz, 1H), 5.72 (d, J = 5.5 Hz, 1H), 5.59 (dd, J = 8.2, 2.0 Hz, 1H), 5.36 (d, J = 5.6 Hz, 1H), 5.21 (s, 1H), 4.21 (dd, J = 12.0, 3.3 Hz, 1H), 4.14 (dd, J = 12.0, 5.1 Hz, 1H), 4.02–3.97 (m, 1H), 3.94–3.88 (m, 2H), 2.58 (hept, J = 7.0 Hz, 1H), 1.11 (dd, J = 7.0, 1.0 Hz, 6H). ¹H and ¹³C NMR matched the previously reported values.⁵

80 g Scale. 5'-O-Isobutyryl cytidine (80 g, 255 mmol, 1.0 equiv) was dissolved in 70% aq 1-butanol (20 V) and then hydroxylamine sulfate (134 g, 817 mmol, 3.2 equiv) was added. The mixture was stirred vigorously and heated to 75–80 °C for 40 h. The mixture was cooled to room temperature, the layers were then separated, and 1-butanol was distilled from the organic layer yielding a solid white, crude material (85 g). This crude material was then dissolved in water (2 V) and heated to 60–65 °C for 60 min. After completely dissolved, the mixture was allowed to cool to room temperature and stirred for 1 h and then cooled to 5–10 °C

and stirred for 3 h. The solid was then filtered and wet-washed with MTBE (3 V) to obtain 56.0 g of molnupiravir. The purification was repeated again for a final yield of 50.0 g (58% corrected) with 97% purity (qNMR).

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomega.1c00772>.

Route cost calculations, additional experimental details for the transformations of compounds **8** to **10** and **10** to **7**, NMR data for compounds **14**, **11**, **9**, **10**, and **7** (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Timothy F. Jamison – Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States; orcid.org/0000-0002-8601-7799; Email: tfj@mit.edu

Authors

Grace P. Ahlqvist – Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

Catherine P. McGeough – Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

Chris Senanayake – TCG GreenChem, Inc., Process R&D Center at Princeton South, Ewing, New Jersey 08628, United States

Joseph D. Armstrong – TCG GreenChem, Inc., Process R&D Center at Princeton South, Ewing, New Jersey 08628, United States

Ajay Yadaw – TCG GreenChem, Inc., Process R&D Center at Princeton South, Ewing, New Jersey 08628, United States

Sarabindu Roy – TCG GreenChem, Inc., Process R&D Center at Princeton South, Ewing, New Jersey 08628, United States

Saeed Ahmad – Medicines for All Institute, Richmond, Virginia 23298, United States

David R. Snead – Medicines for All Institute, Richmond, Virginia 23298, United States; orcid.org/0000-0003-1239-533X

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acsomega.1c00772>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors thank the Bill and Melinda Gates Foundation for their longstanding support of our research. This work was supported by the National Science Foundation Graduate Research Fellowship under Grant No. 1745302 (G.P.A.).

■ REFERENCES

- (1) Cross, R. Merck & Co. Joins Race for COVID-19 Vaccines and Therapies. *Chem. Eng. News* **2020**, *98*, 12.
- (2) Sheahan, T. P.; Sims, A. C.; Zhou, S.; Graham, R. L.; Pruijssers, A. J.; Agostini, M. L.; Leist, S. R.; Schäfer, A.; Dinnon, K. H.; Stevens, L. J.; Chappell, J. D.; Lu, X.; Hughes, T. M.; George, A. S.; Hill, C. S.; Montgomery, S. A.; Brown, A. J.; Bluemling, G. R.; Natchus, M. G.; Saindane, M.; Kolykhalov, A. A.; Painter, G.; Harcourt, J.; Tamin, A.; Thornburg, N. J.; Swanstrom, R.; Denison, M. R.; Baric, R. S. An

orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci. Transl. Med.* **2020**, *12*, No. eabb5883.

(3) Cox, R. M.; Wolf, J. D.; Plemper, R. K. Therapeutically administered ribonucleoside analogue MK-4482/EIDD-2801 blocks SARS-CoV-2 transmission in ferrets. *Nat. Microbiol.* **2021**, *6*, 11–18.

(4) Painter, G. R.; Bluemling, G. R.; Natchus, M. G.; Guthrie, D. N4-Hydroxycytidine and Derivatives and Anti-Viral Uses Related Thereto. WO2019113462A12019

(5) Vasudevan, N.; Ahlqvist, G. P.; McGeough, C. P.; Paymode, D. J.; Cardoso, F. S. P.; Lucas, T.; Dietz, J.-P.; Opatz, T.; Jamison, T. F.; Gupton, F. B.; Snead, D. R. A concise route to MK-4482 (EIDD-2801) from cytidine. *Chem. Commun.* **2020**, *56*, 13363–13364.

(6) Kool, E. T.; Crisalli, P.; Chan, K. M. Fast Alpha Nucleophiles: Structures that Undergo Rapid Hydrazone/Oxime Formation at Neutral pH. *Org. Lett.* **2014**, *16*, 1454–1457.

(7) Alder, C. M.; Hayler, J. D.; Henderson, R. K.; Redman, A. M.; Shukla, L.; Shuster, L. E.; Sneddon, H. F. Updating and further expanding GSK's solvent sustainability guide. *Green Chem.* **2016**, *18*, 3879–3890.

(8) Ortiz, C.; Ferreira, M. L.; Barbosa, O.; Santos, J. C. S.; dos Rodrigues, R. C.; Berenguer-Murcia, A.; Briand, L. E.; Fernandez-Lafuente, R. Novozym 435: the "perfect" lipase immobilized biocatalyst? *Catal. Sci. Technol.* **2019**, *9*, 2380–2420.

(9) (a) Chen, J.-W.; Wu, W.-T. Regeneration of immobilized *Candida antarctica* lipase for transesterification. *J. Biosci. Bioeng.* **2003**, *95*, 466–469. (b) Chowdhury, A.; Mitra, D. A kinetic study on the Novozyme 435-catalyzed esterification of free fatty acids with octanol to produce octyl esters. *Biotechnol. Prog.* **2015**, *31*, 1494–1499. (c) Zhu, S.-G.; Huang, J.-X.; Zhang, G.-M.; Chen, S.-X.; Zhang, F.-L. Development of a Practical Enzymatic Synthesis of Isosorbide-2-acetate. *Org. Process Res. Dev.* **2018**, *22*, 1548–1552. (d) Shin, M.; Seo, J.; Baek, Y.; Lee, T.; Jang, M.; Park, C. Novel and Efficient Synthesis of Phenethyl Formate via Enzymatic Esterification of Formic Acid. *Biomolecules* **2020**, *10*, No. 70.